A Comprehensive *Echinacea* Germplasm Collection Located at the North Central Regional Plant Introduction Station, Ames, Iowa

Joe-Ann H. McCoy, Mark P. Widrlechner, and Jeffrey D. Carstens

USDA/ARS North Central Regional Plant Introduction Station, G212 Agronomy Hall, Iowa State University, Ames, Iowa 50011 U.S.A.

Echinacea is a well-established, high-value crop, both as an ornamental and dietary supplement. A comprehensive collection of Echinacea germplasm is currently held at the USDA-ARS North Central Regional Plant Introduction Station (NCRPIS) in Ames, Iowa, and is available via seed distribution for research purposes http://www.ars-grin.gov/npgs. The NCRPIS's mission includes: (1) The conservation of genetically diverse crop germplasm through collection and acquisition; (2) The conduct of germplasm-related research; and (3) The encouragement of the use of the germplasm collections and associated information for research, crop improvement, and product development.

Representing all nine species collected throughout their respective North American geographic ranges, the *Echinacea* collection includes 159 accessions (Table 1). Extensive morphological characterization data associated with the collection have been assembled and are available to researchers to aid in selection criteria. The collection has been used extensively for various research projects ranging from ornamental breeding studies for the horticulture trade to HPLC analysis of metabolites of interest to the phytopharmaceutical industry.

Germplasm is collected and made available for distribution through a series of steps. Those steps include: (1) Acquisition and exploration; (2) Regeneration and evaluation; (3) Dormancy and germination studies; (4) Seed propagation in field cages with pollinating insects; (5) Harvesting, drying, cleaning, and processing seeds; (6) Long-term storage under controlled conditions; and (7) Distribution for research purposes.

Table 1. Echinacea accessions conserved at the NCRPIS.

Echinacea taxa	Accessions (no.)	Available accessions
E. angustifolia DC.	6	5
E. angustifolia var. angustifolia	39	34
${\it E. angustifolia} \ {\rm var.} \ {\it strigosa} \ {\rm McGregor}$	2	2
E. atrorubens Nutt.	5	5
E. hybrid	2	2
$E.\ laevigata^{\star}$ (F.E. Boynton & Beadle) S.F. Blake	10	*7
E. pallida (Nutt.) Nutt.	45	41
E. paradoxa (Norton) Britton var. neglecta McGre	egor 5	4
E. paradoxa var. paradoxa	5	5
E. purpurea (L.) Moench	17	10
E. sanguinea Nutt.	9	3
$E.\ simulata\ {\it McGregor}$	10	8
E. tennesseensis* (Beadle) Small	4	*4
82% of the collection is available	159	130

^{*}Federally endangered, requires letter of intent or appropriate permit.

This publication was partially supported by grant number P01 ES012020 from the National Institute of Environmental Health Sciences (NIEHS) and the Office of Dietary Supplements (ODS), NIH. "Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIEHS, NIH."