

Back to the future: insights learned over many years— relevant then, now, and for the future[©]

C.E. Whitcomb^a

Lacebark Inc., 2104 N. Cottonwood Road, Stillwater, Oklahoma 74075, USA.

- 1) Roots come first. Watch any seed germinate, from anywhere on the planet and it always sends down a root before producing a shoot. As the root system goes, so goes the plant (Harris et al., 1967; Hathaway and Whitcomb, 1977; Kramer and Kozlowski, 1960; Waisel et al., 2002; Whitcomb, 1981, 1988a, b, 2001).
- 2) Plants run on energy, just like everything else! Focus on energy production improves growth, health and all other aspects (Cobb and Mills, 1988; Gordon and Larson, 1968, 1970; Larson and Gordon, 1969; Whitcomb, 1988a, b).
- 3) Energy produced is not uniformly distributed. The priority of distribution: flowers, fruits, leaves, stems and, finally roots. Any reduction in energy affects roots first (Cobb and Mills, 1988; Gordon and Larson, 1968, 1970; Larson and Gordon, 1969; Whitcomb, 1988a, b).
- 4) Energy produced in plant tops, mostly stays there. Energy produced by mid-section leaves, mostly goes to flowers, fruits and new growth. Most energy going to the root system is produced in leaves on lower branches. Increase in root branching, increases nutrient and water absorption and energy production (Gordon and Larson, 1968, 1970; Reich et al., 1980; Whitcomb and Euchner, 1979; Whitcomb, 1988a, b).
- 5) If in doubt, ask the plant! Try 3 to 7 treatments, use uniform liners, replicate 8 to 10 times and watch. In nearly all cases, plants will tell you their preference. If no clear answer, change treatments or rates and try again. A computer is not needed to do valid research (Whitcomb et al., 1969, 1975; Whitcomb, 1988a, b).
- 6) Your first loss is your best loss! If you purchase 500 liners, only pot up the good ones. You will be saving money by tossing the marginal ones. If in doubt, throw it out! At any stage, culling marginal plants is the wise thing to do (Hathaway and Whitcomb, 1977, 1984; Watson and Himelick, 1982; Whitcomb, 1981, 1988a, b).
- 7) When your pH meter breaks, save your money—do not replace it! Any pH reflects only proportion of acids vs. bases and tells you nothing about “what acids” or “what bases”; pH is a common scapegoat (Lucas and Davis, 1961; Skimina, 1987; Whitcomb, 1985, 1988a, b; Young, 1988).
- 8) I often see the “optimum” pH, but plants show problems. On the other hand, if total nutrition is near optimum, pH will be in the “optimum” range (Lucas and Davis, 1961; Whitcomb et al., 1978; Whitcomb, 1988a, b; Yeager et al., 1983).
- 9) More is not better: especially as it relates to micronutrients. It is NOT how much available iron, but how much iron relative to manganese, relative to boron, relative to copper, etc. All six micronutrients have an associated inner-dependency (Hathaway and Whitcomb, 1984; Whitcomb et al., 1977, 1981; Whitcomb, 1979, 1988a, b).
- 10) If you have what appears to be micronutrient deficiencies or toxicities go back to item 5 and compare plant growth using your current micronutrient source vs. my original Micromax[®]. For example, there are four major sources of iron sulfate, but two work poorly, one fair and one very well. The one that works best also costs more. The same is true for the other micronutrient elements. Cheap is—well, cheap (Whitcomb et al., 1977, 1981; Whitcomb, 1979, 1988a, b).
- 11) When I did a factorial study with the six micronutrient elements (729 treatments), all four test species grew best with the same combination. Forget specialty fertilizers and specialty mixes: get your growth medium, nutrition and drainage right - and you can grow anything in one simple mix (Whitcomb, 1988a, b).
- 12) Just because a plant is native to a location, does not mean that is where it grows best.

^aE-mail: carl@lacebarkinc.com

Ecologically it may be optimal or suboptimal for that environment. Two striking examples; *Pinus radiata* which is native to California, grows far better in Southern Australia and New Zealand. Likewise, *Melaleuca leucadendra*, native to Australia, grows far better in South Florida (Kramer and Kozlowski, 1960; Studer et al., 1978; Waisel et al., 2002; Whitcomb, 1988a, b).

- 13) Work with the plant. Minimize dictating to the plant. Always add Micromax® at 3/4 lb. and 18-6-12 Osmocote (the original 8 to 9 month release, single coating) at 6 lb./cu. yd to the propagation mix for seeds or cuttings. Do not make the plant wait until you decide to provide nutrition (Appleton and Whitcomb, 1983; Carney and Whitcomb, 1983; Tukey and Tukey, 1959; Tukey, 1962; Whitcomb, 1979, 1988a, b).
- 14) Chemistry of irrigation water is the most commonly overlooked factor affecting plant nutrition. Most common problems: excess sodium, high bicarbonates and high calcium. Calcium is the bully in container growth media. Avoid excess calcium (Daughtry, 1988; Good and Tukey, 1966; Rader et al., 1943; Skimina, 1987; Tukey and Tukey, 1959; Tukey, 1962; Whitcomb et al., 1977; Whitcomb, 1985, 1988a, b; Young, 1988).
- 15) Drainable pore space and Darcy's Law (summarized): water will move from a coarse texture to a fine texture readily. Water will not move from a fine texture to a coarse texture until near saturation. Any mix in a container is fine textured relative to the drain holes. Percent drainable pore space should be about 20% (Davis and Whitcomb, 1975; Threadgill, 1983; Whitcomb, 1972, 1988a, b; White and Mastalerz, 1966).
- 16) Pots with vertical slots are just pots that loose water faster. Sidewall openings provide no benefits unless roots are guided into the openings for air-pruning (Whitcomb, 1972, 1988a, b; Whitcomb and Williams, 1983).
- 17) The 4 in. (10 cm) rule. When actively growing root tips are killed by dehydration (air-pruning) or root tip trapping, increased branching occurs along the root axis from about 4 in. back. Place a plant started in an 18 cell RootMaker® tray which is about 4 in. square into a container 10 to 12 in. in diameter, such as a RootMaker® 3- or 5-gal container. Roots grow out, are air-pruned at the sidewall and branch profusely back to the face of the original ball. The resulting fibrous root system exploits the full volume of the container for maximum absorption of water and nutrients (Dickinson and Whitcomb, 1982; Tinus, 1978; Whitcomb, 1988a, b).
- 18) Killing root tips with toxic levels of copper or zinc or ... creates more complications than benefits. Let that one die! (Whitcomb, 1988a, b).
- 19) Any more than 30% shade and you are using shade as a crutch. Light drives the energy production system. (See #4) (Jacobs, 1954; Knox and Hamilton, 1982; Neal, 1969; Neel and Harris, 1971; Telewski and Pruyn, 1998; Whitcomb, 1988a, b).

Know what you know. Know what you do not know. Do not get the two mixed up!

Literature cited

- Appleton, B., and Whitcomb, C.E. (1983). Effects of container size and transplant date on the growth of tree seedlings. *J. Environ. Hortic.* 1, 89-93.
- Carney, M., and Whitcomb, C.E. (1983). Effects of two slow-release fertilizers on the propagation and subsequent growth of three woody plants. *J. Environ. Hortic.* 1, 55-58.
- Cobb, G.S., and Mills, D. (1988). Top pruning and root growth: practical implications. *Comb. Proc. Intl. Plant Prop. Soc.* 38, 470-471.
- Daughtry, W. (1988). Control of *Phytophthora* and *Pythium* by chlorination of irrigation water. *Comb. Proc. Intl. Plant Prop. Soc.* 38, 420-422.
- Davis, R., and Whitcomb, C.E. (1975). Effects of propagation container size on development of high quality tree seedlings. *Comb. Proc. Intl. Plant Prop. Soc.* 25, 251-257.
- Dickinson, S., and Whitcomb, C.E. (1982). Root development of transplanted seedlings in bottomless milk cartons. *J. Arboric.* 8, 223-224.
- Good, G.L., and Tukey, H.B., Jr. (1966). Leaching of metabolites from cuttings propagated under intermittent mist. *J. Am. Soc. Hortic. Sci.* 89, 727-733.

- Gordon, J.C., and Larson, P.R. (1968). The seasonal course of photosynthesis, respiration and distribution of carbon in young *Pinus resinosa* trees as related to wood formation. *Plant Physiol.* 43 (10), 1617–1624 <https://doi.org/10.1104/pp.43.10.1617>. PubMed
- Gordon, J.C., and Larson, P.R. (1970). Redistribution of C-labeled reserve food in young red pines during shoot elongation. *For. Sci.* 16, 14–20.
- Harris, R.W., Long, D., and Davis, W.B. (1967). Root pruning improves nursery tree quality. *J. Am. Soc. Hortic. Sci.* 96, 105–108.
- Hathaway, R.D., and Whitcomb, C.E. (1977). Propagation of *Quercus* seedlings in bottomless containers with Osmocote. *J. Arboric.* 3, 208–212.
- Hathaway, R.D., and Whitcomb, C.E. (1984). Nutrition and performance of container-grown Japanese pine seedlings. *J. Environ. Hortic.* 2, 9–12.
- Jacobs, R.M. (1954). The effects of wind sway on the form and development of *Pinus radiata* D. Don. *Aust. J. Bot.* 2 (1), 35–51 <https://doi.org/10.1071/BT9540035>.
- Knox, G.W., and Hamilton, D.F. (1982). Rooting of *Berberis* and *Ligustrum* cuttings from stock plants grown at selected light intensities. *Sci. Hortic. (Amsterdam)* 16 (1), 85–90 [https://doi.org/10.1016/0304-4238\(82\)90027-9](https://doi.org/10.1016/0304-4238(82)90027-9).
- Kramer, P.J., and Kozlowski, T.T. (1960). *Physiology of Trees* (New York: McGraw-Hill Book Co.).
- Larson, P.R., and Gordon, J.C. (1969). Leaf development, photosynthesis and C₁₄ distribution in *Populus deltoides* seedlings. *Am. J. Bot.* 56 (9), 1058–1066 <https://doi.org/10.1002/j.1537-2197.1969.tb09760.x>.
- Lucas, R.E., and Davis, J.K. (1961). Relationships between pH values of organic soils and availabilities of 12 plant nutrients. *Soil Sci.* 92 (3), 177–182 <https://doi.org/10.1097/00010694-196109000-00005>.
- Neal, P.L. (1969). Growth factors in trunk development of young trees. *Proc. Intl. Shade Tree Conf.* 45, 46–49.
- Neel, P.L., and Harris, R.W. (1971). Motion-induced inhibition of elongation and induction of dormancy in liquidambar. *Science* 173 (3991), 58–59 <https://doi.org/10.1126/science.173.3991.58>. PubMed
- Rader, L.F., Jr., White, L.M., and Whittaker, C.W. (1943). The salt index—a measure of the effect of fertilizers on the concentration of the soil solution. *Soil Sci.* 55 (3), 201–218 <https://doi.org/10.1097/00010694-194303000-00001>.
- Reich, P.B., Jeskey, R.O., Johnson, P.S., and Hinckley, T.M. (1980). Periodic root and shoot growth in oak. *For. Sci.* 26, 590–598.
- Skimina, C. (1987). A 17-year case history of research and implementation of water recycling on container nursery stock. *Comb. Proc. Intl. Plant Prop. Soc.* 37, 82–87.
- Studer, E.J., Steponkus, P.L., Good, G.L., and Wiest, S.C. (1978). Root hardiness of container-grown ornamentals. *HortScience* 13, 172–174.
- Telewski, F.W., and Pruyn, M.L. (1998). Thigmomorphogenesis: a dose response to flexing in *Ulmus americana* seedlings. *Tree Physiol.* 18 (1), 65–68 <https://doi.org/10.1093/treephys/18.1.65>. PubMed
- Threadgill, C.C. (1983). Effects of container depth, diameter, and media on the rooting of cuttings. Masters thesis (Stillwater, OK: Okla. Sta. Univ.).
- Tinus, R.W. (1978). Root system configuration is important to long tree life. *Comb. Proc. Intl. Plant Prop. Soc.* 28, 58–63.
- Tukey, H.B., Jr. (1962). Leaching of metabolites from above-ground plant parts, with special reference to cuttings used for propagation. *Comb. Proc. Intl. Plant Prop. Soc.* 12, 63–70.
- Tukey, H.B., and Tukey, H.B., Jr. (1959). Practical implications of nutrient losses from plant foliage by leaching. *Proc. Am. Soc. Hortic. Sci.* 74, 671–676.
- Waisel, Y., Amramm, E., and Uzi, K. (2002). *Plant Roots: the Hidden Half*, 3rd edn (New York, Basel: Marcel Dekker, Inc.).
- Watson, G.W., and Himelick, E.B. (1982). Root distribution of nursery trees and its relationship to transplanting success. *J. Arboric.* 8, 225–229.
- Whitcomb, C.E. (1972). Growth of *Carissa grandiflora* 'Boxwood Beauty' in varying media, containers, micronutrient levels. *Florida Nurseryman* 17, 12–13, 43.
- Whitcomb, C.E. (1979). Effects of Micromax micronutrients and Osmocote on growth of tree seedlings in containers. *Okla. Agri. Exp. Sta. Res. Rept. P-791*, 42–44.

- Whitcomb, C.E. (1981). Growing Tree Seedlings in Containers. Bulletin #755 (Okla. Agri. Exp. Sta.).
- Whitcomb, C.E. (1985). Water quality and plant production in containers. *Comb. Proc. Intl. Plant Prop. Soc.* 35, 672–677.
- Whitcomb, C.E. (1988a). Calcium, magnesium and irrigation water. *Comb. Proc. Intl. Plant Prop. Soc.* 38, 425–429.
- Whitcomb, C.E. (1988b). *Plant Production in Containers* (Stillwater, Oklahoma: Lacebark Inc.).
- Whitcomb, C.E. (2001). Seedling development: the critical first days. *Comb. Proc. Intl. Plant Prop. Soc.* 51, 610–614.
- Whitcomb, C.E., and Euchner, L.K. (1979). Effects of shade levels on growth of container nursery stock. *Nursery Res. J.* 6, 1–11.
- Whitcomb, C.E., and Williams, J.D. (1983). A stair-step container for improved root growth. *HortScience* 20, 66–67.
- Whitcomb, C.E., Roberts, E.C., and Landers, R.Q. (1969). A connecting pot technique for root competition investigations between woody plants or between woody and herbaceous plants. *Ecology* 50 (2), 326–328 <https://doi.org/10.2307/1934860>.
- Whitcomb, C.E., Rackley, J.G., and Bean, R. (1975). The effects of multiple liners per container on growth and visual grade of woody ornamentals. *Nursery Res. J.* 3, 1–13.
- Whitcomb, C.E., Storjohann, A., and Gibson, J.D. (1977). Effects of time of transplanting container-grown tree seedlings on subsequent growth and development. *Okla. Agri. Exp. Sta. Res. Rept. P-777*, 37–39.
- Whitcomb, C.E., Gibson, J., and Storjohann, A. (1978). Effects of Osmocote formulations in rooting media. *Am. Nurse* 147 (11), 66–68.
- Whitcomb, C.E., Storjohann, A., and Warde, W.D. (1981). Micromax micronutrients for improved plant growth. *Comb. Proc. Intl. Plant Prop. Soc.* 30, 462–467.
- White, J.W., and Mastalerz, J.W. (1966). Soil moisture as related to “container capacity”. *Proc. Am. Soc. Hortic. Sci.* 89, 758–765.
- Yeager, T.H., Wright, R.D., and Donohue, S.J. (1983). Comparison of pour-through and saturated pine bark extract N, P, K, and pH levels. *J. Am. Soc. Hortic. Sci.* 108, 112–114.
- Young, R. (1988). Treating high bicarbonate water. *Comb. Proc. Intl. Plant Prop. Soc.* 38, 423–425.